## WHAT IS CLAIMED IS:

- 1. A polynucleotide fragment comprising a polynucleotide sequence encoding a polypeptide having heparanase catalytic activity, wherein said polypeptide shares at least 70% homology with SEQ ID NOs:10, 14 or 44, as determined using default parameters of a DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the University of Wisconsin.
- 2. The polynucleotide fragment of claim 1, wherein said polynucleotide sequence includes nucleotides 63-1691 of SEQ ID NO:9.
- 3. The polynucleotide fragment of claim 1, wherein said polynucleotide sequence includes nucleotides 63-721 of SEQ ID NO:9.
- 4. The polynucleotide fragment of claim 1, wherein said polynucleotide is as set forth in SEQ ID NO:9.
- 5. The polynucleotide fragment of claim 1, wherein said polynucleotide sequence includes a segment of SEQ ID NO:9, said segment encodes said polypeptide having said heparanase catalytic activity.

- 6. The polynucleotide fragment of claim 1, wherein said polypeptide includes an amino acid sequence as set forth in SEQ ID NOs:10, 14 or 44.
- 7. The polynucleotide fragment of claim 1, wherein said polypeptide includes a segment of SEQ ID NOs:10, 14 or 44 said segment harbors said heparanase catalytic activity.
- 8. The polynucleotide fragment of claim 1, wherein said polynucleotide sequence is selected from the group consisting of double stranded DNA, single stranded DNA and RNA.
- 9. A polynucleotide sequence as set forth in SEQ ID NOs:9, 13, 42 or 43.
- 10. A polynucleotide sequence at least 70% homologous to SEQ ID NOs:9, 13, 42 or 43, as determined using default parameters of a DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the University of Wisconsin, wherein said polynucleotide sequence encodes a polypeptide having heparanase catalytic activity.
- 11. A vector comprising a polynucleotide sequence encoding a polypeptide having heparanase catalytic activity, wherein said polypeptide

shares at least 70% homology with SEQ ID NOs:10, 14 or 44, as determined using default parameters of a DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the University of Wisconsin.

- 12. The vector of claim 11, wherein said polynucleotide sequence includes nucleotides 63-1691 of SEQ ID NO:9.
- 13. The vector of claim 11, wherein said polynucleotide sequence includes nucleotides 63-721 of SEQ ID NO:9.
- 14. The vector of claim 11, wherein said polynucleotide sequence is as set forth in SEQ ID NO:9.
- 15. The vector of claim 11, wherein said polynucleotide sequence includes a segment of SEQ ID NO:9, said segment encodes said polypeptide having said heparanase catalytic activity.
- 16. The vector of claim 11, wherein said polypeptide includes an amino acid sequence as set forth in SEQ ID NOs:10, 14 or 44.

- 17. The vector of claim 11, wherein said polypeptide includes a segment of SEQ ID NOs:10, 14 or 44, said segment harbors said heparanase catalytic activity.
- 18. The vector of claim 11, wherein said polynucleotide sequence is selected from the group consisting of double stranded DNA, single stranded DNA and RNA.
- 19. The vector of claim 11, wherein said vector is a baculovirus vector.
- 20. A host cell comprising an exogenous polynucleotide fragment including a polynucleotide sequence encoding a polypeptide having heparanase catalytic activity, wherein said polypeptide shares at least 70% homology with SEQ ID NOs:10, 14 or 44, as determined using default parameters of a DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the University of Wisconsin.
- 21. The host cell of claim 20, wherein said polynucleotide sequence includes nucleotides 63-1691 of SEQ ID NO:9.

- 22. The host cell of claim 20, wherein said polynucleotide sequence includes nucleotides 63-721 of SEQ ID NO:9.
- 23. The host cell of claim 20, wherein said polynucleotide sequence is as set forth in SEQ ID NO:9.
- 24. The host cell of claim 20, wherein said polynucleotide sequence includes a segment of SEQ ID NO:9, said segment encodes said polypeptide having said heparanase catalytic activity.
- 25. The host cell of claim 20, wherein said polypeptide includes an amino acid sequence as set forth in SEQ ID NOs:10, 14 or 44.
- 26. The host cell of claim 20, wherein said polypeptide includes a segment of SEQ ID NOs:10, 14 or 44 said segment harbors said heparanase catalytic activity.
- 27. The host cell of claim 20, wherein said polynucleotide sequence is selected from the group consisting of double stranded DNA, single stranded DNA and RNA.

- 28. A host cell expressing a recombinant heparanase, wherein said recombinant heparanase shares at least 70% homology with SEQ ID NOs:10, 14 or 44, as determined using default parameter of a DNA sequence analysis software package developed by the Genetic Computer (Group (GCG) at the University of Wisconsin.
- 29. A heparanase overexpression system comprising a cell overexpressing heparanase catalytic activity, wherein said heparanase catalytic activity is effected by a heparanase sharing at least 70% homology with SEQ ID NOs:10, 14 or 44, as determined using default parameters of a DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the University of Wisconsin.
  - 30. The host cell of claim 20, wherein said cell is an insect cell.